

Hazardous

& Toxic

Food Guide for Dogs

Seek Immediate Veterinary attention/advice

FATTY FOODS AND FAT TRIMMINGS

Such as Chicken Wings, Pizza, Burgers, Ribs etc.

Upset stomach, nausea, vomiting, pancreatitis

RAW SALMON

Fresh caught, freshwater fish carrying infected parasites can poision dogs if eaten raw Yomiting, fever, diarrhea, weakness, swollen lymph nodes

RAW MEAT & FISH

Can contain salmonella and E. coli

Vomiting, fever, enlarged lymph nodes

RAW EGGS

May contaion harmful bacteria and risk of salmonella toxicity. Long-term ingestion of raw egg whites can lead to biotin deficiency.

BONES



Cooked bones can splinter, causing internal lacerations, obstruction, choking

Vomiting, lethargy, diarrhea, abdominal discomfort, drooling

NUTS

Almonds, Pecans, Pistachios, Walnuts, Macadamia Nuts, Hickory Nuts

Upset stomach, Pancreatitis *Moldy Walnuts can cause vomiting, tremors and seizures 🔨

GARLIC, ONIO, SHALLOTS, LEEKS, CHIVES

Poisonous to dogs, Toxic in large amounts. Toxic doses of garlic can cause damage to the red blood cells leading to anemia. Signs of garlic poisoning can be delayed and not apparent for several days.

Lethargy, pale gums, increased heart rate, increased respiratory rate, weakness, exercise intolerance, and collapse 🤼

MUSHROOMS 🔷

Some varieties of wild mushrooms are toxic and may cause severe clinical signs or even death. Some mushroom toxins will affect pets very rapidly (within 15-30 minutes of ingestion) while others will not produce signs for many hours

Vomiting, diarrhea, ataxia, tremors, seizures, disorientation, black-tarry stools, inappropriate thirst or urination, Liver or Kidney failure 🔼

XYLITOL

Sugar substitute used in gum, breath mints, toothpaste, candy, some peanut butter, pudding, jam, children's vitamins, nasal spray.

Vomiting, diarrhea, tremors, hyperactivity, seizures, lifethreatening low blood sugar, Liver failure

CHOCOLATE

Chocolate toxicity varies greatly depending on the type and amount of chocolate ingested. The darker and less sweet the chocolate, the more toxic it can be to dogs. Baker's chocolate and dark chocolate pose the greatest risk of toxicity, while white chocolate carries the lowest risk.

Ingestion in toxic amounts can result in vomiting, diarrhea, hyperactivity, elevated heart rate, abnormal heart rhythm, tremors, seizures, and possibly even death (though death is rare in pets who receive prompt and appropriate treatment)



Hazardous

& Toxic

Food Guide for Dogs

Seek Immediate
Veterinary
attention/advice

MILK AND LACTOSE PRODUCTS Dogs lack the enzyme lactase which is needed to digest lactose

can Cause

Upset stomach, diarrhea

GRAPES, RAISINS, SULTANS & CURRENTS

Toxic even in small amounts. Can cause renal (kidney) failure in Sympton

Vomiting, lethargy, diarrhea, abdominal pain, anorexia, and acute renal failure

RHUBARB -

Leaves are Toxic

Drooling, vomiting, diarrhea, lethargy, weakness, tremors, bloody urine, and changes in thirst and urination.

APRICOTS,
PEACHES,
PLUMS,
CHERRY
TREE/SHRUB
BLACK/CHOKE
CHERRY

All parts of these plants other than the ripe pulp around the seeds are considered toxic and contain cyanide.

Symptoms

Dilated pupils, difficulty breathing, inadequate oxygen levels, bright red gums, shock, and death can be seen.

ALCOHOL,
UNBAKED
YEAST DOUGH,
DESSERTS
WITH ALCOHOL

Can cause alcohol poisioning. Alcohol can cause dangerous drops in blood sugar, blood pressure, and body temperature. Severely intoxicated animals can potentially experience seizures and respiratory failure.

Drooling, vomiting,

depression, lethargy,
incoordination, retching,
weakness, collapse, decreased
respiratory rate, seizures

Used in home brewing of Beer. Toxic to dogs.

Signs of hop poisoning include malignant hyperthermia (with temperatures exceeding 105°F/40.6°C). Death has been reported in dogs poisoned by hops within 6 hours of ingestion (without treatment)

Hyperthermia, increased breathing, racing heart rate, anxiety, vomiting, abnormal clotting, and even death

CAFFEINE:
COFFEE
GROUNDS,
TEA BAGS,
ENERGY
DRINKS,
DIET PILLS

Toxicity can vary depending on the amount ingested. In severe cases, seizures, collapse and death are possible.

Symptoms

Panting, hyperactivity, vomiting, elevated heart rate, high blood pressure, abnormal heart rhythms, tremors, and elevated body temperature may be seen.

Commonly used to make holiday ornaments. Usually contains high salt content. Salt toxicity can happen quickly and can result in seizures and death.

Vomiting, diarrhea, lethargy, incoordination, excessive thirst or urination. In severe cases, tremors, seizures, coma, and even death are possible.